

GFM17AF Recombinant Mouse IL-2 (Animal-Free)

Description

Interleukin-2 (IL-2) is an immunomodulatory cytokine that is produced by lymphocytes. IL-2 signals through the IL-2R receptor to induce activated T cell proliferation and promote T cell differentiation. IL-2 also stimulates the proliferation and differentiation of B cells, Natural Killer cells, monocytes, and macrophages.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

Length	150 aa
Molecular Weight	17.4 kDa
Source	E. coli
Accession Number	P04351
Purity	≥95% determined by reducing and non-reducing SDS-PAGE



Specifications

Alternative Names	Interleukin-2, interleukin 2, IL2, IL 2, T cell growth factor, TCGF, Aldesleukin
Biological Activity	Mouse IL-2 is fully biologically active when compared to standard. The activity is determined by the ability to induce CTLL-2 cells proliferation and it is typically less than 5 ng/ml. This corresponds to an expected specific activity of 2×10^5 units/mg.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium citrate, pH 4.0
AA Sequence	MAPTSSSTSS STAEAQQQQQ QQQQQQQHLE QLLMDLQELL SRMENYRNLK LPRMLTFKFY LPKQATELKD LQCLEDELGP LRHVLDLTQS KSFQLEDAEN FISNIRVTVV KLKGSNDTFE CQFDDSATV VDFLRRWIAF CQSIISTSPQ

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile 10 mM acetic acid at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced proliferation of CTLL-2 cells for Mouse IL-2. Cell proliferation was measured to calculate the ED50, which is as expected less than 5 ng/ml.

Reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue, 1 μg of protein was loaded. Mouse IL-2 has a predicted Mw of 17.4 kDa.